



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Multi-trajectory analysis of concurrent changes in physical activity and body mass index among 66,852 public sector employees over 16-year follow-up

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-057692
Article Type:	Original research
Date Submitted by the Author:	27-Sep-2021
Complete List of Authors:	Tiusanen, Roosa; TYKS Turku University Hospital, Department of Physical and Rehabilitation Medicine Saltychev, Mikhail; TYKS Turku University Hospital, Department of Physical and Rehabilitation Medicine Ervasti, Jenni; Finnish Institute of Occupational Health Kivimäki, Mika; University of Helsinki Faculty of Medicine; University College London, Dep. of Epidemiology and Public Health Pentti, Jaana; University of Helsinki, Clinicum, Faculty of Medicine; University of Turku, Department of Public Health Stenholm, Sari; University of Turku, Department of Public Health; University of Turku, Centre for Population Health Research Vahtera, Jussi; University of Turku, Department of Public Health; University of Turku, Centre for Population Health Research
Keywords:	EPIDEMIOLOGY, PREVENTIVE MEDICINE, PUBLIC HEALTH

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Multi-trajectory analysis of concurrent changes in physical activity and body mass index among 66,852 public sector employees over 16-year follow-up**

Running head: **Trajectories of physical activity and BMI**

Roosa Tiusanen<sup>1</sup>, Mikhail Saltychev<sup>1</sup>, Jenni Ervasti<sup>2</sup>, Mika Kivimäki<sup>2,3,4</sup>, Jaana Pentti<sup>3,5,6</sup>, Sari Stenholm<sup>5,6\*</sup>, Jussi Vahtera<sup>5,6\*</sup>

<sup>1</sup> Department of Physical and Rehabilitation Medicine, Turku University Hospital and University of Turku, Turku, Finland

<sup>2</sup> Finnish Institute of Occupational Health, Helsinki, Finland

<sup>3</sup> Clinicum, Faculty of Medicine, University of Helsinki, Helsinki, Finland

<sup>4</sup> Department of Epidemiology and Public Health, University College London Medical School, London, United Kingdom

<sup>5</sup> Department of Public Health, University of Turku and Turku University Hospital, Turku, Finland

<sup>6</sup> Centre for Population Health Research, University of Turku and Turku University Hospital; Turku, Finland

\* Shared senior authorship

*Address for correspondence:*

Roosa Tiusanen, Department of Physical and Rehabilitation Medicine, Turku University Hospital, PO Box 52, FI-20521, Turku, Finland

E-mail: roemti@utu.fi, Tel.: +358 40 8363878



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**DECLARATIONS**

*Ethics approval and consent to participate:*

The ethics committee of the Hospital District of Helsinki and Uusimaa approved the study (registration number HUS/1210/2016). Consent to participate was obtained from each participant.

*Consent for publication:*

Not applicable

*Availability of data and materials:*

We are allowed to share anonymised questionnaire data of the Finnish Public Sector Study by application for with bona fide researchers with an established scientific record and bona fide organisations. For information about the Finnish Public Sector Study contact Prof. Mika Kivimaki mika.kivimaki[at]helsinki.fi / Dr. Jenni Ervasti jenni.ervasti[at]ttl.fi.

*Competing interests:*

The authors declare that they have no competing interests.

*Funding:*

This study was supported by funding granted by the Academy of Finland (Grants 332030 to SS; 633666 to MK; 321409 and 329240 to JV); NordForsk (to MK and JV); the UK MRC (Grant K013351 to MK); Hospital District of Southwest Finland (to SS)

*Author contributions*

All the authors (RT, MS, JE, MK, JP, SS, JV) substantially contributed to the conception and design of the work, the interpretation of the results and revising it critically for important intellectual content. JE, JV and MK were responsible for the acquisition of data for the work. MS and JP were responsible for the statistical analysis. RT and MS were responsible for drafting the work. All the authors (RT,

MS, JE, MK, JP, SS, JV) have finally approved the version to be published and they are agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Acknowledgements:*

None to declare

*Word count:* 2566

*References:* 22

### **Study importance**

*What is already known about this subject:* Physical activity decreases and BMI increases with advancing age when studied separately.

*What this study adds:* Physical activity and BMI are interconnected and there is a negative association between physical activity and BMI. In this study, four clusters of BMI and physical activity trajectories were identified.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

58     **ABSTRACT**

59     **Background**

60     Physical inactivity and body weight tend to increase with age, but their concurrent changes across life

61     is less studied. The objective was to identify concurrent developmental trajectories of physical activity

62     and Body Mass Index (BMI) over 16 years.

63     **Methods**

64     66,852 participants from the Finnish Public Sector cohort study were included. Physical activity during

65     leisure time and commuting and self-reported BMI were assessed with questionnaires. Participants

66     were divided into two age groups; ≤50 and >50 years at baseline. Group-based multi-trajectory

67     modeling was used to investigate the clusters of concurrent BMI and physical activity trajectories over

68     16 years. For sensitivity analysis, the respondents were stratified by gender.

69     **Results**

70     At baseline, mean age was 44.7 (SD 9.4) years, BMI 25.1 (SD 4.1) kg/m<sup>2</sup> and physical activity 27.7 (SD

71     24.8) MET-h/week. Four clusters of concurrent BMI and physical activity trajectories were identified

72     in both age groups. Overall, there was negative association between BMI and physical activity at

73     baseline with BMI increasing and physical activity decreasing over time. The decline in physical activity

74     and increase in BMI were steeper among the respondents with obesity or severe obesity, who had

75     moderately low or low level of physical activity.

76     **Conclusions**

77     Changes in BMI and physical activity were interconnected.

78

79     **Keywords**

80 "Physical activity"; "Body Mass Index"; "Population Dynamics"[Mesh]; "Population  
81 Characteristics"[Mesh]; "Population Health"[Mesh]; ("Prevalence"[Mesh]); "Longitudinal  
82 Studies"[Mesh]

83

#### 84 **Strengths and limitations of this study**

85 Long follow-up of 16 years

86 Large sample size of 66,852 participants

87 Repeated measurements of physical activity and BMI

88 Only leisure-time physical activity was taken into account, leaving out work-related activity

89 BMI and physical activity were self-reported, which may cause bias

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**INTRODUCTION**

Both obesity and physical inactivity have negative impact on multiple aspects of health and they increase the risk of mortality <sup>1-3</sup>. Ageing is associated with gaining weight and decreasing physical activity <sup>4-6</sup>, but less is known whether these changes occur simultaneously and how much heterogeneity there is in the developmental trajectories of body weight and physical activity.

Few studies have examined heterogeneity in weight development over time more closely. A study amongst 30-year-old US war veterans identified five different, but all increasing, trajectories of body mass index (BMI) over 6-year follow-up <sup>6</sup>. However, the steepness of trajectories varied: while the non-obese participants showed only a small increase in BMI, the increase was much steeper among the participants with obesity. Another study from the US conducted on 60-year-old overweight participants identified seven weight trajectories of which most showed either stable overweight, continuously increasing BMI or relapse after weight loss. Even in the two trajectory groups showing decrease in BMI the participants remained overweight.<sup>7</sup>.

Physical activity has also been reported to change over time. Leisure time physical activity among women has previously been reported to increase until age of 50 years and start to decrease after that.<sup>4</sup> For men, the change in leisure time physical activity has been reported to vary between different types of activity - while moderate physical activity increased, low and high levels decreased.<sup>5</sup> Studies concerning trajectories of physical activity have found variation in development of activity. A 22-year follow-up study from Canada among initially 18 to 60 year-olds has identified trajectories of consistently inactive, increasing, consistently active and decreasing leisure time physical activity.<sup>8</sup> Another study conducted in the US among 120 initially overweight people aged 54 (±9) years has measured activity with pedometers and identified "sedentary" and "low active" groups (decreasing daily count of steps), "somewhat active" group (persistent daily count of steps) and "active" group (increased daily count of steps) in 18-month follow-up.<sup>9</sup>

The association between higher levels of physical activity and lower BMI has been established in adults <sup>10 11</sup>, and there has been some evidence that this association might be most pronounced

when physical activity exceeds 150 min/week.<sup>10</sup> There is, however, limited knowledge on simultaneous changes in these two factors. In short-term follow-up (18 months) among overweight 54-year-old Canadians, a trajectory with increasing activity has been associated with a trajectory of greater weight loss.<sup>9</sup> There is yet little knowledge on these two factors over longer follow-up. It is also unknown whether developmental patterns of BMI and physical activity differ by age or by gender.

To address the gap in the literature, the objective of this study was to examine concurrent changes in BMI and physical activity over 16-year follow-up by using a group-based multi-trajectory analysis. While conventional statistics show a trajectory of average change of outcome over time, group-based trajectory modeling can distinguish and describe subpopulations (clusters), which may differ substantially from each other and from the average trajectory seen in the entire population. The aim was also to examine, whether the distinguished trajectories are different for less than 50-year-olds and over 50-year-olds and whether the results are different when the study population is stratified by gender.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**METHODS**

**Study population**

Participants were drawn from the Finnish Public Sector (FPS) cohort study of employees of 10 towns and six hospital districts. Data included responses to five questionnaire surveys administered to the FPS sub-cohorts in four-year intervals from 2000 to 2017 (average response rate 70%). The baseline was the response given in 2000 or 2004. Participants who had reported their BMI and physical activity in at least two waves were included in the analysis. The ethics committee of the Hospital District of Helsinki and Uusimaa approved the study (registration number HUS/1210/2016).

Physical activity was assessed with a questionnaire at all four-year intervals. The respondents were asked to estimate their average weekly hours of leisure-time physical activity/exercise and commuting activity within the previous year. The time spent on activity at each intensity level in hours per week was multiplied by the average energy expenditure of each activity, expressed in metabolic equivalent of task (MET).<sup>12</sup> The MET is a ratio of rate of energy expenditure reflecting the amount of consumed energy compared to resting. One MET unit of 3.5 ml of oxygen per kg per minute corresponds to oxygen consumption when calmly sitting down.

The BMI was defined as weight/height<sup>2</sup> (kg/m<sup>2</sup>) based on self-reported body weight and height. The interpretation of the mean level of BMI trajectories was based on the following categorization: normal weight (<25 kg/m<sup>2</sup>), overweight (25 to 29.9 kg/m<sup>2</sup>), obese (30-34.9 kg/m<sup>2</sup>) and severely obese (≥35 kg/m<sup>2</sup>). Of the respondents, only 934 (1%) had BMI ≤18.5, and thus, for the matter of clarity, BMI <25 was considered “normal”. Age was defined in full years.

**Statistical analysis**

The characteristics of participants were reported as means and standard deviations or as absolute numbers and percentage when appropriate.

Group-based multi-trajectory analysis (GBTA) was used to distinguish different developmental trajectories for physical activity and BMI, both treated as continuous variables. A censored (known also as 'regular') normal model of group-based multi-trajectory analysis was used. The goodness of model fit was judged by running the procedure several times with a number of subpopulations starting from one up to six. The Bayesian Information Criterion (BIC), Akaike information criterion (AIC) and average posterior probability (APP) were used as criteria to confirm the goodness of fit. A cubic regression was applied. The cut-off for the smallest group was set at  $\geq 5\%$ . The trajectory analysis was conducted on two age-groups  $\leq 50$  and  $> 50$  years as previous studies have suggested that changes in BMI and physical activity may vary depending on the age<sup>13 14</sup>. The sensitivity analysis was conducted by dividing both age groups by gender.

All the analyses were performed using Stata/IC Statistical Software: Release 16. College Station (StataCorp LP, TX, USA). The additional Stata module 'traj' was required to conduct group-based trajectory analysis. The module is freely available for both SAS® and Stata software (Jones and Nagin 1999; 2013).

#### **Patient and public involvement**

Participants of research were not involved in setting the study question and outcome measures and were not involved in the design and implementation of the study or writing the manuscript.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**RESULTS**

During the 16-year follow-up, the 66,852 participants had reported body weight and height on average in 3.5 (SD 1.3) study waves and physical activity in 3.6 (SD 1.3) study waves. The sample was predominated by 53,468 women (80%). In the younger group ( $\leq 50$ -year-olds) mean age was 39.8 (SD 7.2), BMI at baseline was 24.6 (SD 4.0)  $\text{kg/m}^2$  and average physical activity was 28.8. (SD 25.5) MET-h/week. In the older group ( $>50$ -year-olds), age was 55.0 (SD 2.9), BMI 25.6 (SD 4.2)  $\text{kg/m}^2$  and physical activity 26.7 (SD 24.1) MET-h/week.

A four-trajectory model was chosen as the five-trajectory model had resulted in a smallest group below a pre-agreed cut-off of 5% (Table 1). Four concurrent trajectories of BMI and physical activity were identified for both age groups (Figure 1 and Figure 2):

- Group 1 (38% among  $< 50$  years, 32% among  $\geq 50$  years): Individuals with normal weight (BMI  $<25 \text{ kg/m}^2$ ) and high level of physical activity (30-35 MET-h).
- Group 2 (39% among  $< 50$  years, 42% among  $\geq 50$  years): Individuals with overweight (BMI 25 to 30  $\text{kg/m}^2$ ) and moderately high level of physical activity (25-30 MET-h).
- Group 3 (18% among  $< 50$  years, 21% among  $\geq 50$  years): Individuals with obesity (BMI 30 to 35  $\text{kg/m}^2$ ) and moderately low level of physical activity (20-25 MET-h).
- Group 4 (5% among  $< 50$  years, 5% among  $\geq 50$  years): Individuals with severe obesity (BMI  $>35 \text{ kg/m}^2$ ) and low level of physical activity ( $<20$  MET-h).

**Group 1: Individuals with normal weight and high level of physical activity**

In this group, the younger respondents demonstrated a stable high level of physical activity with a slight rise towards the end of follow-up and their BMI increased slightly throughout the follow-up. For the older respondents, the level of physical activity decreased markedly during the follow-up, even if there was a slight rising pattern in the middle of follow-up. At the same time, the trajectory of BMI remained flat.

**Group 2: Individuals with overweight and moderately high level of physical activity**

1  
2  
3 195 In this group, the level of physical activity declined in both age groups, but the decline was steeper  
4  
5 196 among the older respondents. In younger respondents, the decrease of physical activity slowed down  
6  
7 197 slightly towards the end of follow-up. Simultaneously, BMI was steadily growing among younger  
8  
9 198 respondents, while remaining relatively flat in older group.

11  
12 199 **Group 3: Individuals with obesity and moderately low level of physical activity**

13  
14 200 The physical activity and BMI trajectories were similar to the trajectories observed in group of  
15  
16 201 overweight individuals with moderately high level of physical activity (group #2), but with a slightly  
17  
18 202 steeper decline in physical activity and steeper increase in BMI.

19  
20 203 **Group 4: Individuals with severe obesity and low level of physical activity**

21  
22 204 Also in this group, physical activity decreased and BMI increased. In younger respondents, this  
23  
24 205 development slowed down at the end follow-up for both physical activity and BMI. Instead, in older  
25  
26 206 respondents, the decrease in physical activity accelerated towards the end of follow-up with  
27  
28 207 simultaneous slight decline in BMI.

29  
30 208 **Sensitivity analysis**

31  
32 209 Stratifying the respondents by gender in addition to age resulted in similar findings with few  
33  
34 210 exceptions (supplementary figures E1-E4). Among normal weight or overweight respondents, the  
35  
36 211 decline in physical activity was steeper among men compared to women.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**DISCUSSION**

This prospective cohort study in 66,852 public sector employees followed repeatedly by 4-year intervals investigated trajectories of concurrent changes in BMI and physical activity over 16 years. Four trajectory clusters were identified for both participants aged  $\leq 50$  and for those  $> 50$  years: 1) individuals with normal weight and high level of physical activity; 2) individuals with overweight and moderately high level of physical activity; 3) individuals with obesity and moderately low level of physical activity; and 4) individuals with severe obesity and low level of physical activity. On average, BMI increased and physical activity decreased during the follow-up. Some trajectories demonstrated, however, distinctive features. Over time, the respondents with normal weight or overweight gained only a little weight while preserved a high or moderately high level of physical activity, even if the intensity of physical activity mildly decreased especially in older respondents. The decrease in physical activity and increase in BMI were steeper among the respondents with obesity or severe obesity, who had moderately low or low level of physical activity already at the start of the follow-up. Among the normal weight or overweight respondents, decline in physical activity was steeper among men compared to women.

The observed age-related weight gain is in line with previous studies<sup>4-6 15</sup>, as well as the decline in physical activity<sup>4 16 17</sup>. Previous studies have also shown that an increase in BMI slows down with advancing age, and this was also supported by the present findings – the rise in BMI was steeper in the younger respondents<sup>18 19</sup>. During the follow-up, the decline in physical activity mirrored the increase in BMI. Similar findings have been reported before – several studies conducted among middle-aged adults have observed an association between physical activity and weight gain<sup>10 11 20 21</sup>. This association has been described to be dose-dependent – physically active individuals gain less weight than inactive peers.<sup>11</sup> Current results support this finding, since the increase in BMI was less steep in the more active groups. The amount of activity needed to prevent weight gain has been debated. Some studies have concluded that current activity recommendations are not sufficient enough to prevent weight gain and that there is a need for higher activity to remain in the normally

weighted category<sup>10 11 20</sup>. This is in line with the current findings – only high physical activity was associated with normal weight.

The strengths of the study were long follow-up of 16 years, repeated measurements on physical activity and BMI, and a large sample size. For our knowledge, there are no previous multi-trajectory analyses of the relation between physical activity and BMI conducted in adults.

The study has also some limitations. Physical activity was self-reported and only leisure-time and commuting activity were inquired. Thus, physical activity at work was not considered. The distribution of physical activity intensity was skewed – most of the participants were at least somewhat active, and even in the least active group the mean activity level was approximately 18 MET/week. BMI was also based on self-reported weight and height, which may cause recall and information bias, possibly resulting in under-reporting of body weight<sup>22</sup>. Most of the participants had BMI above 25 indicating overweight or obesity (62% in the age group of  $\leq 50$  years and 68% in the older), which may reflect the current overweight and obesity pandemic. The cohort included predominantly working-age women employed in public sector. Therefore, the results might not be directly reflected on the entire population, since there might be variation in behavior, for instance among unemployed people or entrepreneurs. Moreover, a public sector often employs people with higher socioeconomic status, who might have more knowledge and financial resources to healthy lifestyle choices compared to manual workers.

## Conclusions

Changes in BMI and physical activity were interconnected. The normal weight or overweight respondents gained only a little weight while preserved a high or moderately high level of physical activity. Compared to normal weight trajectories, the decrease of physical activity and increase in BMI were markedly steeper among the obese or severely obese trajectories, who also had moderately low or low level of physical activity. Among the normal weight and overweight trajectories, decline in physical activity was steeper among men compared to women. Since physical inactivity and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

263     overweight are both risk factors for many diseases, more research is needed to develop interventions  
264     that could simultaneously affect both.

For peer review only

265 **LIST OF ABBREVIATIONS**

266 BMI: Body Mass Index

267 FPS: Finnish Public Sector cohort study

268 MET: Metabolic Equivalent of Task

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

269 REFERENCES

270 1. Bull F, Goenka S, Lambert V, et al. Physical Activity for the Prevention of Cardiometabolic Disease.  
271 In: rd, Prabhakaran D, Anand S, et al., eds. Cardiovascular, Respiratory, and Related  
272 Disorders. Washington (DC)2017.

273 2. Bigaard J, Frederiksen K, Tjonneland A, et al. Waist circumference and body composition in  
274 relation to all-cause mortality in middle-aged men and women. *Int J Obes (Lond)*  
275 2005;29(7):778-84. doi: 10.1038/sj.ijo.0802976 [published Online First: 2005/05/27]

276 3. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World*  
277 *Health Organ Tech Rep Ser* 2000;894:i-xii, 1-253. [published Online First: 2001/03/10]

278 4. Sui X, Zhang J, Lee DC, et al. Physical activity/fitness peaks during perimenopause and BMI change  
279 patterns are not associated with baseline activity/fitness in women: a longitudinal study with  
280 a median 7-year follow-up. *Br J Sports Med* 2013;47(2):77-82. doi: 10.1136/bjsports-2011-  
281 090888 [published Online First: 2012/07/10]

282 5. Droyvold WB, Holmen J, Midthjell K, et al. BMI change and leisure time physical activity (LTPA): an  
283 11-y follow-up study in apparently healthy men aged 20-69 y with normal weight at  
284 baseline. *Int J Obes Relat Metab Disord* 2004;28(3):410-7. doi: 10.1038/sj.ijo.0802569  
285 [published Online First: 2004/01/16]

286 6. Rosenberger PH, Ning Y, Brandt C, et al. BMI trajectory groups in veterans of the Iraq and  
287 Afghanistan wars. *Prev Med* 2011;53(3):149-54. doi: 10.1016/j.ypmed.2011.07.001  
288 [published Online First: 2011/07/21]

289 7. Fitzpatrick SL, Rosales AG, Brown SD, et al. Behavioural and psychosocial factors associated with 5-  
290 year weight trajectories within the PORTAL Overweight/Obesity Cohort. *Obes Sci Pract*  
291 2020;6(3):272-81. doi: 10.1002/osp4.411 [published Online First: 2020/06/12]

292 8. Barnett TA, Gauvin L, Craig CL, et al. Distinct trajectories of leisure time physical activity and  
293 predictors of trajectory class membership: a 22 year cohort study. *Int J Behav Nutr Phys Act*  
294 2008;5:57. doi: 10.1186/1479-5868-5-57 [published Online First: 2008/11/08]

295 9. Imes CC, Zheng Y, Mendez DD, et al. Group-Based Trajectory Analysis of Physical Activity Change  
296 in a US Weight Loss Intervention. *J Phys Act Health* 2018;15(11):840-46. doi:  
297 10.1123/jpah.2017-0484 [published Online First: 2018/10/14]

298 10. Jakicic JM, Powell KE, Campbell WW, et al. Physical Activity and the Prevention of Weight Gain in  
299 Adults: A Systematic Review. *Med Sci Sports Exerc* 2019;51(6):1262-69. doi:  
300 10.1249/MSS.0000000000001938 [published Online First: 2019/05/17]

301 11. Moholdt T, Wisloff U, Lydersen S, et al. Current physical activity guidelines for health are  
302 insufficient to mitigate long-term weight gain: more data in the fitness versus fatness debate  
303 (The HUNT study, Norway). *Br J Sports Med* 2014;48(20):1489-96. doi: 10.1136/bjsports-  
304 2014-093416 [published Online First: 2014/05/02]

305 12. Leskinen T, Stenholm S, Aalto V, et al. Physical activity level as a predictor of healthy and chronic  
306 disease-free life expectancy between ages 50 and 75. *Age Ageing* 2018;47(3):423-29. doi:  
307 10.1093/ageing/afy016 [published Online First: 2018/03/17]

308 13. Stenholm S, Pulakka A, Kawachi I, et al. Changes in physical activity during transition to  
309 retirement: a cohort study. *Int J Behav Nutr Phys Act* 2016;13:51. doi: 10.1186/s12966-016-  
310 0375-9 [published Online First: 2016/04/17]

311 14. Stenholm S, Solovieva S, Viikari-Juntura E, et al. Change in body mass index during transition to  
312 statutory retirement: an occupational cohort study. *Int J Behav Nutr Phys Act* 2017;14(1):85.  
313 doi: 10.1186/s12966-017-0539-2 [published Online First: 2017/06/28]

314 15. Yang Y, Dugue PA, Lynch BM, et al. Trajectories of body mass index in adulthood and all-cause  
315 and cause-specific mortality in the Melbourne Collaborative Cohort Study. *BMJ Open*  
316 2019;9(8):e030078. doi: 10.1136/bmjopen-2019-030078 [published Online First:  
317 2019/08/12]

16. Achttien RJ, van Lieshout J, Wensing M, et al. The decline in physical activity in aging people is not modified by gender or the presence of cardiovascular disease. *Eur J Public Health* 2020;30(2):333-39. doi: 10.1093/eurpub/ckz159 [published Online First: 2019/09/29]
17. McAuley E, Hall KS, Motl RW, et al. Trajectory of declines in physical activity in community-dwelling older women: social cognitive influences. *J Gerontol B Psychol Sci Soc Sci* 2009;64(5):543-50. doi: 10.1093/geronb/gbp049 [published Online First: 2009/06/17]
18. Brown WJ, Kabor E, Clark BK, et al. Maintaining a Healthy BMI: Data From a 16-Year Study of Young Australian Women. *Am J Prev Med* 2016;51(6):e165-e78. doi: 10.1016/j.amepre.2016.09.007 [published Online First: 2016/11/22]
19. Lewis CE, Jacobs DR, Jr., McCreath H, et al. Weight gain continues in the 1990s: 10-year trends in weight and overweight from the CARDIA study. Coronary Artery Risk Development in Young Adults. *Am J Epidemiol* 2000;151(12):1172-81. doi: 10.1093/oxfordjournals.aje.a010167 [published Online First: 2000/07/25]
20. Erlichman J, Kerbey AL, James WP. Physical activity and its impact on health outcomes. Paper 2: Prevention of unhealthy weight gain and obesity by physical activity: an analysis of the evidence. *Obes Rev* 2002;3(4):273-87. doi: 10.1046/j.1467-789x.2002.00078.x [published Online First: 2002/12/03]
21. Hankinson AL, Daviglius ML, Bouchard C, et al. Maintaining a high physical activity level over 20 years and weight gain. *JAMA* 2010;304(23):2603-10. doi: 10.1001/jama.2010.1843 [published Online First: 2010/12/16]
22. Connor Gorber S, Tremblay M, Moher D, et al. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev* 2007;8(4):307-26. doi: 10.1111/j.1467-789X.2007.00347.x [published Online First: 2007/06/21]



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

344   **TABLES**

345   Table 1. Goodness of fit of group-based trajectory analysis models. The chosen models are shown in  
346   bold.

Model	Smallest group		BIC <sup>1</sup>	AIC <sup>2</sup>	APP <sup>3</sup>
	n	%			
<51 years					
1-cluster	31,797	100%	-905,561	-905,509	1
2-cluster	8,234	26%	-869,531	-869,432	0.94
3-cluster	3,331	10%	-851,542	-851,397	0.92
<b>4-cluster</b>	<b>1,490</b>	<b>5%</b>	<b>-841,703</b>	<b>-841,510</b>	<b>0.89</b>
5-cluster	898	3%	-835,396	-835,157	0.87
>50 years					
1-cluster	35,055	100%	-869,200	-869,148	1
2-cluster	9,690	28%	-836,174	-836,076	0.93
3-cluster	3,845	11%	-819,600	-819,454	0.91
<b>4-cluster</b>	<b>1,888</b>	<b>5%</b>	<b>-809,601</b>	<b>-809,409</b>	<b>0.89</b>
5-cluster	999	3%	-803,977	-803,738	0.87

347   <sup>1</sup> BIC = Bayesian Information Criterion, <sup>2</sup> AIC = Akaike information criterion, <sup>3</sup> APP = Smallest average  
348   posterior probability




Figure 1. Trajectories of physical activity and body mass index amongst respondents <50 years. 95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.

700x510mm (120 x 120 DPI)




Figure 2. Trajectories of physical activity and body mass index amongst respondents > 50 years95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.

700x510mm (120 x 120 DPI)

Table E1. Goodness of fit of group-based trajectory analysis models. The chosen models are shown in bold.

Model	Smallest group		BIC <sup>1</sup>	AIC <sup>2</sup>	APP <sup>3</sup>
	n	%			
Men <51 years					
1-cluster	5,894	100%	-156,412	-156,369	1
2-cluster	1,469	25%	-151,020	-150,938	0.93
3-cluster	509	9%	-148,201	-148,080	0.91
<b>4-cluster</b>	<b>292</b>	<b>5%</b>	<b>-146,715</b>	<b>-146,555</b>	<b>0.88</b>
5-cluster	147	2%	-145,799	-145,600	0.86
Men >50 years					
1-cluster	7,490	100%	-177,574	-177,530	1
2-cluster	1,894	25%	-171,451	-171,368	0.92
3-cluster	622	8%	-168,332	-168,209	0.90
<b>4-cluster</b>	<b>334</b>	<b>4%</b>	<b>-166,442</b>	<b>-166,280</b>	<b>0.88</b>
5-cluster	174	2%	-165,267	-165,066	0.87
Women <51 years					
1-cluster	25,903	100%	-746,837	-746,786	1
2-cluster	6,530	25%	-715,572	-715,475	0.95
3-cluster	2,773	11%	-700,393	-700,250	0.92
<b>4-cluster</b>	<b>1,173</b>	<b>5%</b>	<b>-692,029</b>	<b>-691,840</b>	<b>0.90</b>
5-cluster	745	3%	-686,684	-686,449	0.87
Women >50 years					
1-cluster	27,565	100%	-690,012	-689,961	1
2-cluster	7,608	28%	-662,602	-662,506	0.94
3-cluster	3,164	11%	-649,085	-648,944	0.91
<b>4-cluster</b>	<b>1,536</b>	<b>6%</b>	<b>-641,136</b>	<b>-640,949</b>	<b>0.89</b>
5-cluster	842	3%	-636,666	-636,433	0.86

<sup>1</sup> BIC = Bayesian Information Criterion, <sup>2</sup> AIC = Akaike information criterion, <sup>3</sup> APP = Smallest average posterior probability




Figure E1. Trajectories of physical activity and body mass index (BMI) amongst men < 50 years  
95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.




Figure E2. Trajectories of physical activity and body mass index amongst men > 50 years

95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.




Figure E3. Trajectories of physical activity and body mass index amongst women < 50 years  
95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.




Figure E4. Trajectories of physical activity and body mass index amongst women > 50 years  
95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.



STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page No
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	4
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	6,7
Objectives	3	State specific objectives, including any prespecified hypotheses	7
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	8
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (b) For matched studies, give matching criteria and number of exposed and unexposed	8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	8
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	8
Bias	9	Describe any efforts to address potential sources of bias	9
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	9
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, explain how loss to follow-up was addressed (e) Describe any sensitivity analyses	9 9 9 9 9
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	10
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) Summarise follow-up time (eg, average and total amount)	10
Outcome data	15*	Report numbers of outcome events or summary measures over time	10,11

Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	10,11
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10,11
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	12
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	13
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	12,13
Generalisability	21	Discuss the generalisability (external validity) of the study results	13
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	2

\*Give information separately for exposed and unexposed groups.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at <http://www.strobe-statement.org>.

# BMJ Open

## Concurrent changes in physical activity and body mass index among 66,852 public sector employees over 16-year follow-up: multi-trajectory analysis of a cohort study in Finland

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-057692.R1
Article Type:	Original research
Date Submitted by the Author:	19-Dec-2021
Complete List of Authors:	Tiusanen, Roosa; TYKS Turku University Hospital, Department of Physical and Rehabilitation Medicine Saltychev, Mikhail; TYKS Turku University Hospital, Department of Physical and Rehabilitation Medicine Ervasti, Jenni; Finnish Institute of Occupational Health Kivimäki, Mika; University of Helsinki Faculty of Medicine; University College London, Dep. of Epidemiology and Public Health Pentti, Jaana; University of Helsinki, Clinicum, Faculty of Medicine; University of Turku, Department of Public Health Stenholm, Sari; University of Turku, Department of Public Health; University of Turku, Centre for Population Health Research Vahtera, Jussi; University of Turku, Department of Public Health; University of Turku, Centre for Population Health Research
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Epidemiology
Keywords:	PUBLIC HEALTH, EPIDEMIOLOGY, QUALITATIVE RESEARCH

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Concurrent changes in physical activity and body mass index among 66,852 public sector employees over 16-year follow-up: multi-trajectory analysis of a cohort study in Finland**

Running head: **Trajectories of physical activity and BMI**

Roosa Tiusanen<sup>1</sup>, Mikhail Saltychev<sup>1</sup>, Jenni Ervasti<sup>2</sup>, Mika Kivimäki<sup>2,3,4</sup>, Jaana Pentti<sup>3,5,6</sup>, Sari Stenholm<sup>5,6\*</sup>, Jussi Vahtera<sup>5,6\*</sup>

<sup>1</sup> Department of Physical and Rehabilitation Medicine, Turku University Hospital and University of Turku, Turku, Finland

<sup>2</sup> Finnish Institute of Occupational Health, Helsinki, Finland

<sup>3</sup> Clinicum, Faculty of Medicine, University of Helsinki, Helsinki, Finland

<sup>4</sup> Department of Epidemiology and Public Health, University College London Medical School, London, United Kingdom

<sup>5</sup> Department of Public Health, University of Turku and Turku University Hospital, Turku, Finland

<sup>6</sup> Centre for Population Health Research, University of Turku and Turku University Hospital; Turku, Finland

\* Shared senior authorship

*Address for correspondence:*

Roosa Tiusanen, Department of Physical and Rehabilitation Medicine, Turku University Hospital, PO Box 52, FI-20521, Turku, Finland

E-mail: roemti@utu.fi, Tel.: +358 40 8363878

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**DECLARATIONS**

*Ethics approval and consent to participate:*

The ethics committee of the Hospital District of Helsinki and Uusimaa approved the study (registration number HUS/1210/2016). Written informed consent to participate was obtained from each participant.

*Consent for publication:*

Not applicable

*Data availability statement:*

We are allowed to share anonymised questionnaire data of the Finnish Public Sector Study by application for with bona fide researchers with an established scientific record and bona fide organisations. For information about the Finnish Public Sector Study contact Prof. Mika Kivimaki mika.kivimaki[at]helsinki.fi / Dr. Jenni Ervasti jenni.ervasti[at]ttl.fi.

*Competing interests:*

The authors declare that they have no competing interests.

*Funding:*

This study was supported by funding granted by the Academy of Finland (Grants 332030 to SS; 633666 to MK; 321409 and 329240 to JV); NordForsk (to MK and JV); the UK MRC (Grant K013351 to MK); Hospital District of Southwest Finland (to SS)

*Author contributions*

All the authors (RT, MS, JE, MK, JP, SS, JV) substantially contributed to the conception and design of the work, the interpretation of the results and revising it critically for important intellectual content. JE, JV and MK were responsible for the acquisition of data for the work. MS and JP were responsible

for the statistical analysis. RT and MS were responsible for drafting the work. All the authors (RT, MS, JE, MK, JP, SS, JV) have finally approved the version to be published and they are agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Acknowledgements:*

None to declare

*Word count:* 2496

*References:* 26

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**ABSTRACT**

**Objectives:** To identify concurrent developmental trajectories of physical activity and body mass Index (BMI) over time.

**Design:** Prospective cohort study, repeated survey.

**Setting:** University research department and national institute of occupational health.

**Participants:** 66,852 public sector employees, who have been followed for 16 years.

**Primary and secondary outcome measures:** Shapes of trajectories of changes in physical activity and BMI.

**Results** At baseline, mean age was 44.7 (SD 9.4) years, BMI 25.1 (SD 4.1) kg/m<sup>2</sup> and physical activity 27.7 (SD 24.8) MET-h/week. Four clusters of concurrent BMI and physical activity trajectories were identified. There was negative association between BMI and physical activity.

**Conclusions:** Changes in BMI and physical activity might be interconnected. Decline in physical activity and increase in BMI were steeper among obese respondents with low level of physical activity.

**Trial registration:** The ethics committee of the Hospital District of Helsinki and Uusimaa approved the study (registration number HUS/1210/2016).

**Keywords**

“Physical activity”; “Body Mass Index”; “Population Dynamics”[Mesh]; “Population Characteristics”[Mesh]; “Population Health”[Mesh]; (“Prevalence”[Mesh]); “Longitudinal Studies”[Mesh]

**Article Summary**

**Strengths and limitations of this study**

- Large cohort of 66,852 participants
- Repeated measures of physical activity and BMI during 16 years
- Only leisure-time physical activity was taken into account leaving out work-related activity
- Self-reported nature of estimates of BMI and physical activity might lead to information bias



## INTRODUCTION

Both obesity and physical inactivity have negative impact on multiple aspects of health and they increase the risk of mortality<sup>1-3</sup>. Ageing is associated with gaining weight and decreasing physical activity<sup>4-6</sup>, but less is known whether these changes occur simultaneously and how much heterogeneity there is in the developmental trajectories of body weight and physical activity.

Few studies have examined heterogeneity in weight development over time more closely. A study amongst 30-year-old US war veterans identified five different, but all increasing, trajectories of body mass index (BMI) over 6-year follow-up<sup>6</sup>. However, the steepness of trajectories varied: while the non-obese participants showed only a small increase in BMI, the increase was much steeper among the participants with obesity. Another study from the US conducted on 60-year-old overweight participants identified seven weight trajectories of which most showed either stable overweight, continuously increasing BMI or relapse after weight loss. Even in the two trajectory groups showing decrease in BMI the participants remained overweight.<sup>7</sup>

Physical activity has also been reported to change over time. Leisure time physical activity among women has previously been reported to increase until age of 50 years and start to decrease after that.<sup>4</sup> For men, the change in leisure time physical activity has been reported to vary between different types of activity - while moderate physical activity increased, low and high levels decreased.<sup>5</sup> Studies concerning trajectories of physical activity have found variation in development of activity. A 22-year follow-up study from Canada among initially 18 to 60 year-olds has identified trajectories of consistently inactive, increasing, consistently active and decreasing leisure time physical activity.<sup>8</sup> Another study conducted in the US among 120 initially overweight people aged 54 ( $\pm 9$ ) years has measured activity with pedometers and identified "sedentary" and "low active" groups (decreasing daily count of steps), "somewhat active" group (persistent daily count of steps) and "active" group (increased daily count of steps) in 18-month follow-up.<sup>9</sup>

The association between higher levels of physical activity and lower BMI has been established in adults<sup>10 11</sup>, and there has been some evidence that this association might be most pronounced

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

when physical activity exceeds 150 min/week.<sup>10</sup> There is, however, limited knowledge on simultaneous changes in these two factors. In short-term follow-up (18 months) among overweight 54-year-old Canadians, a trajectory with increasing activity has been associated with a trajectory of greater weight loss.<sup>9</sup> There is yet little knowledge on these two factors over longer follow-up. It is also unknown whether developmental patterns of BMI and physical activity differ by age or by gender.

To address the gap in the literature, the objective of this study was to examine concurrent changes in BMI and physical activity over 16-year follow-up by using a group-based multi-trajectory analysis. While conventional statistics show a trajectory of average change of outcome over time, group-based trajectory modeling can distinguish and describe subpopulations (clusters), which may differ substantially from each other and from the average trajectory seen in the entire population. The aim was also to examine, whether the distinguished trajectories are different for less than 50-year-olds and over 50-year-olds and whether the results are different when the study population is stratified by gender.

## 115 METHODS

### 116 Study population

117 Participants were drawn from the Finnish Public Sector (FPS) cohort study of employees of 10 towns  
 118 and six hospital districts. The FPS has been described in detail elsewhere.<sup>12 13</sup> Data included responses  
 119 to five questionnaire surveys administered to the FPS sub-cohorts in four-year intervals from 2000 to  
 120 2017 (average response rate 70%). The baseline was the response given in 2000 or 2004. Participants  
 121 who had reported their BMI and physical activity in at least two waves were included in the analysis.  
 122 The ethics committee of the Hospital District of Helsinki and Uusimaa approved the study (registration  
 123 number HUS/1210/2016).

124 Physical activity was assessed with a questionnaire at all four-year intervals. The respondents  
 125 were asked to estimate their average weekly hours of leisure-time physical activity/exercise and  
 126 commuting activity within the previous year. The time spent on activity at each intensity level in hours  
 127 per week was multiplied by the average energy expenditure of each activity, expressed in metabolic  
 128 equivalent of task (MET).<sup>14</sup> The MET is a ratio of rate of energy expenditure reflecting the amount of  
 129 consumed energy compared to resting. One MET unit of 3.5 ml of oxygen per kg per minute  
 130 corresponds to oxygen consumption when calmly sitting down. Weekly physical activity was expressed  
 131 as MET-h/week and categorized as low (<14 MET-h/week), moderate (14 to <30 MET-h/week) or high  
 132 ( $\geq 30$  MET-h/week) physical activity levels.<sup>15 16</sup> The definition of physical activity in the survey is  
 133 presented in table 1. As recommended by the American Heart Association, 150 minutes of moderate-  
 134 intensity aerobic exercise each week is needed for optimal cardiovascular health. That is equal to  
 135 about 8.3 MET-h/week.

136 The BMI was defined as weight/height<sup>2</sup> (kg/m<sup>2</sup>) based on self-reported body weight and height. The  
 137 interpretation of the mean level of BMI trajectories was based on the following categorization: normal  
 138 weight (<25 kg/m<sup>2</sup>), overweight (25 to 29.9 kg/m<sup>2</sup>), obese (30-34.9 kg/m<sup>2</sup>) and severely obese ( $\geq 35$

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

kg/m<sup>2</sup>). Of the respondents, only 934 (1%) had BMI  $\leq$ 18.5, and thus, for the matter of clarity, BMI <25 was considered “normal”. Age was defined in full years. The cohort was divided in two approximately even age groups:  $\leq$ 50 (n=31,797, 48%) and >50 years (n=35,055, 52%).

**Statistical analysis**

The characteristics of participants were reported as means and standard deviations or as absolute numbers and percentage when appropriate.

Group-based multi-trajectory analysis (GBTA) was used to distinguish different developmental trajectories for physical activity and BMI, both treated as continuous variables. A censored (known also as ‘regular’) normal model of group-based multi-trajectory analysis was used. The goodness of model fit was judged by running the procedure several times with a number of subpopulations starting from one up to six. The Bayesian Information Criterion (BIC), Akaike information criterion (AIC) and average posterior probability (APP) were used as criteria to confirm the goodness of fit. A cubic regression was applied. The cut-off for the smallest group was set at  $\geq$ 5%. The trajectory analysis was conducted on two age-groups  $\leq$ 50 and >50 years as previous studies have suggested that changes in BMI and physical activity may vary depending on the age<sup>17 18</sup>. The sensitivity analysis was conducted by dividing both age groups by gender. No adjustments for co-variables were made.

All the analyses were performed using Stata/IC Statistical Software: Release 16. College Station (StataCorp LP, TX, USA). The additional Stata module ‘traj’ was required to conduct group-based trajectory analysis. The module is freely available for both SAS® and Stata software (Jones and Nagin 1999; 2013).

**Patient and public involvement**

Participants of research were not involved in setting the study question and outcome measures and were not involved in the design and implementation of the study or writing the manuscript.

## 163 RESULTS

164 During the 16-year follow-up, the 66,852 participants had reported body weight and height on average  
 165 in 3.5 (SD 1.3) study waves and physical activity in 3.6 (SD 1.3) study waves. The sample was  
 166 predominated by 53,468 women (80%). In the younger group ( $\leq 50$ -year-olds) mean age was 39.8 (SD  
 167 7.2), BMI at baseline was 24.6 (SD 4.0)  $\text{kg/m}^2$  and average physical activity was 28.8. (SD 25.5) MET-  
 168 h/week. In the older group ( $>50$ -year-olds), age was 55.0 (SD 2.9), BMI 25.6 (SD 4.2)  $\text{kg/m}^2$  and physical  
 169 activity 26.7 (SD 24.1) MET-h/week.

170 A four-trajectory model was chosen as the five-trajectory model had resulted in a smallest group  
 171 below a pre-agreed cut-off of 5% (Table 2). Four concurrent trajectories of BMI and physical activity  
 172 were identified for both age groups (Figure 1 and Figure 2):

1731. Group 1 (38% among  $\leq 50$  years, 32% among  $> 50$  years): Individuals with normal weight (BMI  $<25$   
 174  $\text{kg/m}^2$ ) and high level of physical activity (30-35 MET-h/week).

1752. Group 2 (39% among  $\leq 50$  years, 42% among  $> 50$  years): Individuals with overweight (BMI 25 to 30  
 176  $\text{kg/m}^2$ ) and moderately high level of physical activity (25-30 MET-h/week).

1773. Group 3 (18% among  $\leq 50$  years, 21% among  $> 50$  years): Individuals with obesity (BMI 30 to 35  $\text{kg/m}^2$ )  
 178 and moderately low level of physical activity (20-25 MET-h/week).

1794. Group 4 (5% among  $\leq 50$  years, 5% among  $> 50$  years): Individuals with severe obesity (BMI  $>35$   $\text{kg/m}^2$ )  
 180 and low level of physical activity ( $<20$  MET-h/week).

### 181 **Group 1: Individuals with normal weight and high level of physical activity**

182 In this group, the younger respondents demonstrated a stable high level of physical activity with a  
 183 slight rise towards the end of follow-up and their BMI increased slightly throughout the follow-up. For  
 184 the older respondents, the level of physical activity decreased markedly during the follow-up, even if  
 185 there was a slight rising pattern in the middle of follow-up. At the same time, the trajectory of BMI  
 186 remained flat.

### 187 **Group 2: Individuals with overweight and moderately high level of physical activity**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

In this group, the level of physical activity declined in both age groups, but the decline was steeper among the older respondents. In younger respondents, the decrease of physical activity slowed down slightly towards the end of follow-up. Simultaneously, BMI was steadily growing among younger respondents, while remaining relatively flat in older group.

**Group 3: Individuals with obesity and moderately low level of physical activity**

The physical activity and BMI trajectories were similar to the trajectories observed in group of overweight individuals with moderately high level of physical activity (group #2), but with a slightly steeper decline in physical activity and steeper increase in BMI.

**Group 4: Individuals with severe obesity and low level of physical activity**

Also in this group, physical activity decreased and BMI increased. In younger respondents, this development slowed down at the end follow-up for both physical activity and BMI. Instead, in older respondents, the decrease in physical activity accelerated towards the end of follow-up with simultaneous slight decline in BMI.

**Sensitivity analysis**

Stratifying the respondents by gender in addition to age resulted in similar findings with few exceptions (supplementary figures E1-E4 and supplementary table E1). Among normal weight or overweight respondents, the decline in physical activity was steeper among men compared to women.

## DISCUSSION

This prospective cohort study in 66,852 public sector employees followed repeatedly by 4-year intervals investigated trajectories of concurrent changes in BMI and physical activity over 16 years. Four trajectory clusters were identified for both participants aged  $\leq 50$  and for those  $> 50$  years: 1) individuals with normal weight and high level of physical activity; 2) individuals with overweight and moderately high level of physical activity; 3) individuals with obesity and moderately low level of physical activity; and 4) individuals with severe obesity and low level of physical activity. On average, BMI increased and physical activity decreased during the follow-up. Some trajectories demonstrated, however, distinctive features. Over time, the respondents with normal weight or overweight gained only a little weight while preserved a high or moderately high level of physical activity, even if the intensity of physical activity mildly decreased especially in older respondents. The decrease in physical activity and increase in BMI were steeper among the respondents with obesity or severe obesity, who had moderately low or low level of physical activity already at the start of the follow-up. Among the normal weight or overweight respondents, decline in physical activity was steeper among men compared to women.

The observed age-related weight gain is in line with previous studies<sup>4-6 19</sup>, as well as the decline in physical activity<sup>4 20 21</sup>. Previous studies have also shown that an increase in BMI slows down with advancing age, and this was also supported by the present findings – the rise in BMI was steeper in the younger respondents<sup>22 23</sup>. During the follow-up, the decline in physical activity mirrored the increase in BMI. Similar findings have been reported before – several studies conducted among middle-aged adults have observed an association between physical activity and weight gain<sup>10 11 24 25</sup>. This association has been described to be dose-dependent – physically active individuals gain less weight than inactive peers.<sup>11</sup> Current results support this finding, since the increase in BMI was less steep in the more active groups. The amount of activity needed to prevent weight gain has been debated. Some studies have concluded that current activity recommendations are not sufficient enough to prevent weight gain and that there is a need for higher activity to remain in the normally

1  
2  
3 231 weighted category <sup>10 11 24</sup>. This is in line with the current findings – only high physical activity was  
4  
5 232 associated with normal weight.  
6

7 233 The strengths of the study were long follow-up of 16 years, repeated measurements on  
8  
9  
10 234 physical activity and BMI, and a large sample size. For our knowledge, there are no previous multi-  
11  
12 235 trajectory analyses of the relation between physical activity and BMI conducted in adults.  
13

14 236 The study has also some limitations. Physical activity was self-reported and only leisure-time  
15  
16 237 and commuting activity were inquired. Thus, physical activity at work was not considered. The  
17  
18 238 distribution of physical activity intensity was skewed – most of the participants were at least  
19  
20 239 somewhat active, and even in the least active group the mean activity level was approximately 18  
21  
22 240 MET-h/week, which is approximately the equivalent of three hours of brisk walking weekly. BMI was  
23  
24 241 also based on self-reported weight and height, which may cause recall and information bias, possibly  
25  
26 242 resulting in under-reporting of body weight <sup>26</sup>. Most of the participants had BMI above 25 indicating  
27  
28 243 overweight or obesity (62% in the age group of  $\leq 50$  years and 68% in the older), which may reflect the  
29  
30 244 current overweight and obesity pandemic. The cohort included predominantly working-age women  
31  
32 245 employed in public sector. Therefore, the results might not be directly reflected on the entire  
33  
34 246 population, since there might be variation in behavior, for instance among unemployed people or  
35  
36 247 entrepreneurs. Moreover, a public sector often employs people with higher socioeconomic status,  
37  
38 248 who might have more knowledge and financial resources to healthy lifestyle choices compared to  
39  
40 249 manual workers.  
41  
42  
43  
44

45 250 The results may be of interest for both clinicians and stockholders when applying measures  
46  
47 251 targeting increasing physical activity and controlling weight especially among people of middle-age.  
48  
49 252 Additionally, the information on the established trajectories may give people more motivation to  
50  
51 253 change their health behavior. Further research may reveal risk factors that affect developmental  
52  
53 254 trajectories seen in this study. Such factors may be, for example, gender, socio-economic status,  
54  
55 255 smoking, alcohol consumption and concurrent health disorders among others.  
56  
57  
58

59 256 **Conclusions**  
60



257 Changes in BMI and physical activity might be interconnected. The normal weight or overweight  
258 respondents gained only a little weight while preserved a high or moderately high level of physical  
259 activity. Compared to normal weight trajectories, the decrease of physical activity and increase in BMI  
260 were markedly steeper among the obese or severely obese trajectories, who also had moderately low  
261 or low level of physical activity. The findings were similar for both age groups. Among the normal  
262 weight and overweight trajectories, decline in physical activity was steeper among men compared to  
263 women. Since physical inactivity and overweight are both risk factors for many diseases, more  
264 research is needed to develop interventions that could simultaneously affect both.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**LIST OF ABBREVIATIONS**

- BMI: Body Mass Index
- FPS: Finnish Public Sector cohort study
- MET: Metabolic Equivalent of Task

**FIGURE LEGENDS**

- Figure 1. Trajectories of physical activity and body mass index amongst respondents  $\leq 50$  years
- 95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.
- Figure 2. Trajectories of physical activity and body mass index amongst respondents  $> 50$  years
- 95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.

## REFERENCES

1. Bull F, Goenka S, Lambert V, et al. Physical Activity for the Prevention of Cardiometabolic Disease. In: rd, Prabhakaran D, Anand S, et al., eds. Cardiovascular, Respiratory, and Related Disorders. Washington (DC)2017.
2. Bigaard J, Frederiksen K, Tjonneland A, et al. Waist circumference and body composition in relation to all-cause mortality in middle-aged men and women. *Int J Obes (Lond)* 2005;29(7):778-84. doi: 10.1038/sj.ijo.0802976 [published Online First: 2005/05/27]
3. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* 2000;894:i-xii, 1-253. [published Online First: 2001/03/10]
4. Sui X, Zhang J, Lee DC, et al. Physical activity/fitness peaks during perimenopause and BMI change patterns are not associated with baseline activity/fitness in women: a longitudinal study with a median 7-year follow-up. *Br J Sports Med* 2013;47(2):77-82. doi: 10.1136/bjsports-2011-090888 [published Online First: 2012/07/10]
5. Droyvold WB, Holmen J, Midthjell K, et al. BMI change and leisure time physical activity (LTPA): an 11-y follow-up study in apparently healthy men aged 20-69 y with normal weight at baseline. *Int J Obes Relat Metab Disord* 2004;28(3):410-7. doi: 10.1038/sj.ijo.0802569 [published Online First: 2004/01/16]
6. Rosenberger PH, Ning Y, Brandt C, et al. BMI trajectory groups in veterans of the Iraq and Afghanistan wars. *Prev Med* 2011;53(3):149-54. doi: 10.1016/j.ypmed.2011.07.001 [published Online First: 2011/07/21]
7. Fitzpatrick SL, Rosales AG, Brown SD, et al. Behavioural and psychosocial factors associated with 5-year weight trajectories within the PORTAL Overweight/Obesity Cohort. *Obes Sci Pract* 2020;6(3):272-81. doi: 10.1002/osp4.411 [published Online First: 2020/06/12]
8. Barnett TA, Gauvin L, Craig CL, et al. Distinct trajectories of leisure time physical activity and predictors of trajectory class membership: a 22 year cohort study. *Int J Behav Nutr Phys Act* 2008;5:57. doi: 10.1186/1479-5868-5-57 [published Online First: 2008/11/08]
9. Imes CC, Zheng Y, Mendez DD, et al. Group-Based Trajectory Analysis of Physical Activity Change in a US Weight Loss Intervention. *J Phys Act Health* 2018;15(11):840-46. doi: 10.1123/jpah.2017-0484 [published Online First: 2018/10/14]
10. Jakicic JM, Powell KE, Campbell WW, et al. Physical Activity and the Prevention of Weight Gain in Adults: A Systematic Review. *Med Sci Sports Exerc* 2019;51(6):1262-69. doi: 10.1249/MSS.0000000000001938 [published Online First: 2019/05/17]
11. Moholdt T, Wisloff U, Lydersen S, et al. Current physical activity guidelines for health are insufficient to mitigate long-term weight gain: more data in the fitness versus fatness debate (The HUNT study, Norway). *Br J Sports Med* 2014;48(20):1489-96. doi: 10.1136/bjsports-2014-093416 [published Online First: 2014/05/02]
12. Kouvonen A, Kivimaki M, Elovainio M, et al. Effort/reward imbalance and sedentary lifestyle: an observational study in a large occupational cohort. *Occup Environ Med* 2006;63(6):422-7. doi: 10.1136/oem.2005.020974 [published Online First: 2006/02/25]
13. Kouvonen A, Kivimaki M, Virtanen M, et al. Effort-reward imbalance at work and the co-occurrence of lifestyle risk factors: cross-sectional survey in a sample of 36,127 public sector employees. *BMC Public Health* 2006;6:24. doi: 10.1186/1471-2458-6-24 [published Online First: 2006/02/09]
14. Leskinen T, Stenholm S, Aalto V, et al. Physical activity level as a predictor of healthy and chronic disease-free life expectancy between ages 50 and 75. *Age Ageing* 2018;47(3):423-29. doi: 10.1093/ageing/afy016 [published Online First: 2018/03/17]
15. Kujala UM, Kaprio J, Sarna S, et al. Relationship of leisure-time physical activity and mortality: the Finnish twin cohort. *JAMA* 1998;279(6):440-4. doi: 10.1001/jama.279.6.440 [published Online First: 1998/02/18]

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

16. Ainsworth BE, Haskell WL, Leon AS, et al. Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc* 1993;25(1):71-80. doi: 10.1249/00005768-199301000-00011 [published Online First: 1993/01/01]

17. Stenholm S, Pulakka A, Kawachi I, et al. Changes in physical activity during transition to retirement: a cohort study. *Int J Behav Nutr Phys Act* 2016;13:51. doi: 10.1186/s12966-016-0375-9 [published Online First: 2016/04/17]

18. Stenholm S, Solovieva S, Viikari-Juntura E, et al. Change in body mass index during transition to statutory retirement: an occupational cohort study. *Int J Behav Nutr Phys Act* 2017;14(1):85. doi: 10.1186/s12966-017-0539-2 [published Online First: 2017/06/28]

19. Yang Y, Dugue PA, Lynch BM, et al. Trajectories of body mass index in adulthood and all-cause and cause-specific mortality in the Melbourne Collaborative Cohort Study. *BMJ Open* 2019;9(8):e030078. doi: 10.1136/bmjopen-2019-030078 [published Online First: 2019/08/12]

20. Achttien RJ, van Lieshout J, Wensing M, et al. The decline in physical activity in aging people is not modified by gender or the presence of cardiovascular disease. *Eur J Public Health* 2020;30(2):333-39. doi: 10.1093/eurpub/ckz159 [published Online First: 2019/09/29]

21. McAuley E, Hall KS, Motl RW, et al. Trajectory of declines in physical activity in community-dwelling older women: social cognitive influences. *J Gerontol B Psychol Sci Soc Sci* 2009;64(5):543-50. doi: 10.1093/geronb/gbp049 [published Online First: 2009/06/17]

22. Brown WJ, Kabir E, Clark BK, et al. Maintaining a Healthy BMI: Data From a 16-Year Study of Young Australian Women. *Am J Prev Med* 2016;51(6):e165-e78. doi: 10.1016/j.amepre.2016.09.007 [published Online First: 2016/11/22]

23. Lewis CE, Jacobs DR, Jr., McCreath H, et al. Weight gain continues in the 1990s: 10-year trends in weight and overweight from the CARDIA study. Coronary Artery Risk Development in Young Adults. *Am J Epidemiol* 2000;151(12):1172-81. doi: 10.1093/oxfordjournals.aje.a010167 [published Online First: 2000/07/25]

24. Erlichman J, Kerbey AL, James WP. Physical activity and its impact on health outcomes. Paper 2: Prevention of unhealthy weight gain and obesity by physical activity: an analysis of the evidence. *Obes Rev* 2002;3(4):273-87. doi: 10.1046/j.1467-789x.2002.00078.x [published Online First: 2002/12/03]

25. Hankinson AL, Daviglius ML, Bouchard C, et al. Maintaining a high physical activity level over 20 years and weight gain. *JAMA* 2010;304(23):2603-10. doi: 10.1001/jama.2010.1843 [published Online First: 2010/12/16]

26. Connor Gorber S, Tremblay M, Moher D, et al. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev* 2007;8(4):307-26. doi: 10.1111/j.1467-789X.2007.00347.x [published Online First: 2007/06/21]

## 366 TABLES

367 Table 1. Defining the level of physical activity in the survey.

What was the intensity and frequency of your average physical activity (leisure or commuting) during the past year (or since the onset of your disease if the disease had begun less than a year ago)?					
Intensity (Mark all four options)	Amount per week				
	None	<½ hour	1 hour	2 – 3 hours	≥ 4 hours
Normal walking or respective					
Brisk walking or respective					
Light jogging or respective					
Brisk jogging or respective					
The responses were converted into MET units according to a following scheme.					
Intensity (Mark all four options)	MET minutes per week				
	None	<½ hour	1 hour	2 – 3 hours	≥ 4 hours
Normal walking or respective	0	69	138	345	550
Brisk walking or respective	0	99	198	495	792
Light jogging or respective	0	210	420	1050	1680
Brisk jogging or respective	0	240	480	1200	1920









368


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 2. Goodness of fit of group-based trajectory analysis models. The chosen models are shown in bold.


Model	Smallest group		BIC <sup>1</sup>	AIC <sup>2</sup>	APP <sup>3</sup>
	n	%			
≤50 years					
1-cluster model	31,797	100%	-905,561	-905,509	1
2-cluster model	8,234	26%	-869,531	-869,432	0.94
3-cluster model	3,331	10%	-851,542	-851,397	0.92
<b>4-cluster model</b>	<b>1,490</b>	<b>5%</b>	<b>-841,703</b>	<b>-841,510</b>	<b>0.89</b>
5-cluster model	898	3%	-835,396	-835,157	0.87
>50 years					
1-cluster model	35,055	100%	-869,200	-869,148	1
2-cluster model	9,690	28%	-836,174	-836,076	0.93
3-cluster model	3,845	11%	-819,600	-819,454	0.91
<b>4-cluster model</b>	<b>1,888</b>	<b>5%</b>	<b>-809,601</b>	<b>-809,409</b>	<b>0.89</b>
5-cluster model	999	3%	-803,977	-803,738	0.87

<sup>1</sup> BIC = Bayesian Information Criterion, <sup>2</sup> AIC = Akaike information criterion, <sup>3</sup> APP = Smallest average posterior probability









For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

Figure E1. Trajectories of physical activity and body mass index (BMI) amongst men < 50 years

95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.




Figure E2. Trajectories of physical activity and body mass index amongst men > 50 years  
95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.




Figure E3. Trajectories of physical activity and body mass index amongst women < 50 years

95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.




Figure E4. Trajectories of physical activity and body mass index amongst women > 50 years  
95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.

Table E1. Goodness of fit of group-based trajectory analysis models. The chosen models are shown in bold.

Model	Smallest group		BIC <sup>1</sup>	AIC <sup>2</sup>	APP <sup>3</sup>
	n	%			
Men <51 years					
1-cluster	5,894	100%	-156,412	-156,369	1
2-cluster	1,469	25%	-151,020	-150,938	0.93
3-cluster	509	9%	-148,201	-148,080	0.91
<b>4-cluster</b>	<b>292</b>	<b>5%</b>	<b>-146,715</b>	<b>-146,555</b>	<b>0.88</b>
5-cluster	147	2%	-145,799	-145,600	0.86
Men >50 years					
1-cluster	7,490	100%	-177,574	-177,530	1
2-cluster	1,894	25%	-171,451	-171,368	0.92
3-cluster	622	8%	-168,332	-168,209	0.90
<b>4-cluster</b>	<b>334</b>	<b>4%</b>	<b>-166,442</b>	<b>-166,280</b>	<b>0.88</b>
5-cluster	174	2%	-165,267	-165,066	0.87
Women <51 years					
1-cluster	25,903	100%	-746,837	-746,786	1
2-cluster	6,530	25%	-715,572	-715,475	0.95
3-cluster	2,773	11%	-700,393	-700,250	0.92
<b>4-cluster</b>	<b>1,173</b>	<b>5%</b>	<b>-692,029</b>	<b>-691,840</b>	<b>0.90</b>
5-cluster	745	3%	-686,684	-686,449	0.87
Women >50 years					
1-cluster	27,565	100%	-690,012	-689,961	1
2-cluster	7,608	28%	-662,602	-662,506	0.94
3-cluster	3,164	11%	-649,085	-648,944	0.91
<b>4-cluster</b>	<b>1,536</b>	<b>6%</b>	<b>-641,136</b>	<b>-640,949</b>	<b>0.89</b>
5-cluster	842	3%	-636,666	-636,433	0.86

<sup>1</sup> BIC = Bayesian Information Criterion, <sup>2</sup> AIC = Akaike information criterion, <sup>3</sup> APP = Smallest average posterior probability

STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page No
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	4
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	6,7
Objectives	3	State specific objectives, including any prespecified hypotheses	7
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	8
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (b) For matched studies, give matching criteria and number of exposed and unexposed	8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	8
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	8
Bias	9	Describe any efforts to address potential sources of bias	9
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	9
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, explain how loss to follow-up was addressed (e) Describe any sensitivity analyses	9 9 9 9 9
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	10
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) Summarise follow-up time (eg, average and total amount)	10
Outcome data	15*	Report numbers of outcome events or summary measures over time	10,11

1	Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	10,11
2			(b) Report category boundaries when continuous variables were categorized	
3			(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
4	Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10,11
5	<b>Discussion</b>			
6	Key results	18	Summarise key results with reference to study objectives	12
7	Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	13
8	Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	12,13
9	Generalisability	21	Discuss the generalisability (external validity) of the study results	13
10	<b>Other information</b>			
11	Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	2

\*Give information separately for exposed and unexposed groups.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at <http://www.strobe-statement.org>.

# BMJ Open

## Concurrent changes in physical activity and body mass index among 66,852 public sector employees over a 16-year follow-up: multi-trajectory analysis of a cohort study in Finland

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-057692.R2
Article Type:	Original research
Date Submitted by the Author:	25-Jan-2022
Complete List of Authors:	Tiusanen, Roosa; TYKS Turku University Hospital, Department of Physical and Rehabilitation Medicine Saltychev, Mikhail; TYKS Turku University Hospital, Department of Physical and Rehabilitation Medicine Ervasti, Jenni; Finnish Institute of Occupational Health Kivimäki, Mika; University of Helsinki Faculty of Medicine; University College London, Dep. of Epidemiology and Public Health Pentti, Jaana; University of Helsinki, Clinicum, Faculty of Medicine; University of Turku, Department of Public Health Stenholm, Sari; University of Turku, Department of Public Health; University of Turku, Centre for Population Health Research Vahtera, Jussi; University of Turku, Department of Public Health; University of Turku, Centre for Population Health Research
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Epidemiology
Keywords:	PUBLIC HEALTH, EPIDEMIOLOGY, QUALITATIVE RESEARCH

SCHOLARONE™  
Manuscripts





I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Concurrent changes in physical activity and body mass index among 66,852 public sector employees over a 16-year follow-up: multi-trajectory analysis of a cohort study in Finland**

Roosa Tiusanen<sup>1</sup>, Mikhail Saltychev<sup>1</sup>, Jenni Ervasti<sup>2</sup>, Mika Kivimäki<sup>2,3,4</sup>, Jaana Pentti<sup>3,5,6</sup>, Sari Stenholm<sup>5,6\*</sup>, Jussi Vahtera<sup>5,6\*</sup>

<sup>1</sup> Department of Physical and Rehabilitation Medicine, Turku University Hospital and University of Turku, Turku, Finland

<sup>2</sup> Finnish Institute of Occupational Health, Helsinki, Finland

<sup>3</sup> Clinicum, Faculty of Medicine, University of Helsinki, Helsinki, Finland

<sup>4</sup> Department of Epidemiology and Public Health, University College London Medical School, London, United Kingdom

<sup>5</sup> Department of Public Health, University of Turku and Turku University Hospital, Turku, Finland

<sup>6</sup> Centre for Population Health Research, University of Turku and Turku University Hospital; Turku, Finland

\* Shared senior authorship

*Address for correspondence:*

Roosa Tiusanen, Department of Physical and Rehabilitation Medicine, Turku University Hospital, PO Box 52, FI-20521, Turku, Finland

E-mail: roemti@utu.fi, Tel.: +358 40 8363878

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

19   **DECLARATIONS**

20   *Ethics approval and consent to participate:*

21   The ethics committee of the Hospital District of Helsinki and Uusimaa approved the study (registration

22   number HUS/1210/2016). Written informed consent to participate was obtained from each

23   participant.

24   *Consent for publication:*

25   Not applicable

26   *Data availability statement:*

27   We are allowed to share anonymised questionnaire data of the Finnish Public Sector Study by

28   application for with bona fide researchers with an established scientific record and bona fide

29   organisations. For information about the Finnish Public Sector Study contact Prof. Mika Kivimaki

30   mika.kivimaki[at]helsinki.fi / Dr. Jenni Ervasti jenni.ervasti[at]ttl.fi.

31   *Competing interests:*

32   The authors declare that they have no competing interests.

33   *Funding:*

34   This study was supported by funding granted by the Academy of Finland (Grants 332030 to SS; 633666

35   to MK; 321409 and 329240 to JV); NordForsk (to MK and JV); the UK MRC (Grant K013351 to MK);

36   Hospital District of Southwest Finland (to SS)

37   *Author contributions*

38   All the authors (RT, MS, JE, MK, JP, SS, JV) substantially contributed to the conception and design of

39   the work, the interpretation of the results and revising it critically for important intellectual content.

40   JE, JV and MK were responsible for the acquisition of data for the work. MS and JP were responsible

for the statistical analysis. RT and MS were responsible for drafting the work. All the authors (RT, MS, JE, MK, JP, SS, JV) have finally approved the version to be published and they are agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Word count: 2632*

*References: 28*

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**ABSTRACT**

**Objectives:** To identify concurrent developmental trajectories of physical activity and body mass index (BMI) over time.

**Design:** Prospective cohort study, repeated survey.

**Setting:** Cohort study in Finland.

**Participants:** 66,852 public sector employees, who have been followed up for 16 years.

**Outcome measures:** Shapes of trajectories of changes in physical activity and BMI.

**Results:** At baseline, mean age was 44.7 (SD 9.4) years, BMI 25.1 (SD 4.1) kg/m<sup>2</sup> and physical activity 27.7 (SD 24.8) MET-h/week. Four clusters of concurrent BMI and physical activity trajectories were identified: 1) normal weight (BMI <25 kg/m<sup>2</sup>) and high level of physical activity (30-35 MET-h/week), 2) overweight (BMI 25 to 30 kg/m<sup>2</sup>) and moderately high level of physical activity (25-30 MET-h/week), 3) obesity (BMI 30 to 35 kg/m<sup>2</sup>) and moderately low level of physical activity (20-25 MET-h/week) and 4) severe obesity (BMI >35 kg/m<sup>2</sup>) and low level of physical activity (<20 MET-h/week). In general, BMI increased and physical activity decreased during the follow-up. Decline in physical activity and increase in BMI were steeper among obese respondents with low level of physical activity.

**Conclusions:** Changes in BMI and physical activity might be interconnected. The results may be of interest for both clinicians and other stakeholders with respect to informing measures targeting increasing physical activity and controlling weight, especially among middle-aged people. Additionally, the information on the established trajectories may give individuals motivation to change their health behavior.

**Keywords**

“Physical activity”; “Body Mass Index”; “Population Dynamics”[Mesh]; “Population Characteristics”[Mesh]; “Population Health”[Mesh]; (“Prevalence”[Mesh]); “Longitudinal Studies”[Mesh]

**Article Summary**

### Strengths and limitations of this study

- Large cohort of 66,852 participants.
- Repeated measures of physical activity and BMI over 16 years.
- Only leisure-time physical activity was taken into account, leaving out work-related activity.
- The self-reported nature of estimates of BMI and physical activity might lead to information bias.

### INTRODUCTION

Both obesity and physical inactivity have negative impact on multiple aspects of health and they increase the risk of mortality<sup>1-3</sup>. Ageing is associated with gaining weight and decreasing physical activity<sup>4-6</sup>, but less is known whether these changes occur simultaneously and how much heterogeneity there is in the developmental trajectories of body weight and physical activity.

Few studies have examined heterogeneity in weight development over time more closely. A study amongst 30-year-old US war veterans identified five different, but all increasing, trajectories of body mass index (BMI) over 6-year follow-up<sup>6</sup>. However, the steepness of trajectories varied: while the non-obese participants showed only a small increase in BMI, the increase was much steeper among the participants with obesity. Another study from the US conducted on 60-year-old overweight participants identified seven weight trajectories of which most showed either stable overweight, continuously increasing BMI or relapse after weight loss. Even in the two trajectory groups showing decrease in BMI the participants remained overweight.<sup>7</sup>

Physical activity has also been reported to change over time. Leisure time physical activity among women has previously been reported to increase until age of 50 years and start to decrease after that.<sup>4</sup> For men, the change in leisure time physical activity has been reported to vary between different types of activity - while moderate physical activity increased, low and high levels decreased.<sup>5</sup>

1  
2  
3 97 Studies concerning trajectories of physical activity have found variation in development of activity. A  
4  
5 98 22-year follow-up study from Canada among initially 18 to 60 year-olds has identified trajectories of  
6  
7 99 consistently inactive, increasing, consistently active and decreasing leisure time physical activity.<sup>8</sup>  
8  
9  
10 100 Another study conducted in the US among 120 initially overweight people aged 54 ( $\pm 9$ ) years has  
11  
12 101 measured activity with pedometers and identified "sedentary" and "low active" groups (decreasing  
13  
14 102 daily count of steps), "somewhat active" group (persistent daily count of steps) and "active" group  
15  
16 103 (increased daily count of steps) in 18-month follow-up.<sup>9</sup>  
17

18  
19 104 The association between higher levels of physical activity and lower BMI has been established  
20  
21 105 in adults <sup>10 11</sup>, and there has been some evidence that this association might be most pronounced  
22  
23 106 when physical activity exceeds 150 min/week.<sup>10</sup> There is, however, limited knowledge on  
24  
25 107 simultaneous changes in these two factors. In short-term follow-up (18 months) among overweight  
26  
27 108 54-year-old Canadians, a trajectory with increasing activity has been associated with a trajectory of  
28  
29 109 greater weight loss.<sup>9</sup> There is yet little knowledge on these two factors over longer follow-up. It is also  
30  
31 110 unknown whether developmental patterns of BMI and physical activity differ by age or by gender.  
32  
33

34 111 To address the gap in the literature, the objective of this study was to examine concurrent  
35  
36 112 changes in BMI and physical activity over 16-year follow-up by using a group-based multi-trajectory  
37  
38 113 analysis. While conventional statistics show a trajectory of average change of outcome over time,  
39  
40 114 group-based trajectory modeling can distinguish and describe subpopulations (clusters), which may  
41  
42 115 differ substantially from each other and from the average trajectory seen in the entire population. The  
43  
44 116 aim was also to examine, whether the distinguished trajectories are different for less than 50-year-  
45  
46 117 olds and over 50-year-olds and whether the results are different when the study population is  
47  
48 118 stratified by gender.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 119 METHODS

### 120 Study population

121 Participants were drawn from the Finnish Public Sector (FPS) cohort study, a dynamic cohort with  
122 follow-up intervals two to four years initiated from 1998/2000. It consists of employees in the  
123 municipal services of 10 Finnish towns and 21 public hospitals, who had a job contract for a minimum  
124 of 6 months. In year 2000, the most common occupations of the respondents were registered nurse  
125 (23%), teacher (19%), practical nurse (13%) and cleaner (10%). The FPS has been described in more  
126 detail elsewhere.<sup>12 13</sup> Data in the current study included responses to five questionnaire surveys  
127 administered in 2000-2002, 2004-2005, 2008-2009 and 2016-2017 to 2017 (average response rate  
128 70%). The baseline was the response given in 2000-2002 or 2004-2005. Participants who had reported  
129 their BMI and physical activity in at least two waves were included in the analysis. The ethics  
130 committee of the Hospital District of Helsinki and Uusimaa approved the study (registration number  
131 HUS/1210/2016).

132 Physical activity was assessed with a questionnaire at all survey waves. The respondents were  
133 asked to estimate their average weekly hours of leisure-time physical activity/exercise and commuting  
134 activity within the previous year. The time spent on activity at each intensity level in hours per week  
135 was multiplied by the average energy expenditure of each activity, expressed in metabolic equivalent  
136 of task (MET).<sup>14</sup> The MET is a ratio of rate of energy expenditure reflecting the amount of consumed  
137 energy compared to resting. One MET unit of 3.5 ml of oxygen per kg per minute corresponds to  
138 oxygen consumption when calmly sitting down. Weekly physical activity was expressed as MET-  
139 h/week and categorized as low (<14 MET-h/week), moderate (14 to <30 MET-h/week) or high (≥30  
140 MET-h/week) physical activity levels.<sup>15 16</sup> This categorization was chosen since physical activity higher  
141 than 14 MET-h/week has been reported to be associated with cardiovascular disease<sup>17</sup> and the activity  
142 level of 30 MET-h/week has been shown to be needed for weight management<sup>18</sup>. 14 MET-h/week is



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

approximately the equivalent of 140 minutes of brisk walking weekly. The definition of physical activity in the survey is presented in supplementary table E1.

The BMI was defined as weight/height<sup>2</sup> (kg/m<sup>2</sup>) based on self-reported body weight and height. The interpretation of the mean level of BMI trajectories was based on the following categorization: normal weight (<25 kg/m<sup>2</sup>), overweight (25 to 29.9 kg/m<sup>2</sup>), obese (30-34.9 kg/m<sup>2</sup>) and severely obese (≥35 kg/m<sup>2</sup>). Of the respondents, only 934 (1%) had BMI ≤18.5, and thus, for the matter of clarity, BMI <25 was considered “normal”. Age was defined in full years. The cohort was divided in two approximately even age groups: ≤50 (n=31,797, 48%) and >50 years (n=35,055, 52%).

**Statistical analysis**

The characteristics of participants were reported as means and standard deviations or as absolute numbers and percentage when appropriate.

Group-based multi-trajectory analysis (GBTA) was used to distinguish different developmental trajectories for physical activity and BMI, both treated as continuous variables. This method is a form of finite mixture modeling for analyzing longitudinal repeated measures data. While conventional statistics show a trajectory of average change of outcome over time, group-based trajectory modeling is able to distinguish and describe subpopulations (clusters) existing within a studied population. A censored (known also as ‘regular’) normal model of group-based multi-trajectory analysis was used. The goodness of model fit was judged by running the procedure several times with a number of trajectory clusters starting from one up to five, until the smallest group was below the pre-agreed cut-off at ≥5%. The Bayesian Information Criterion (BIC), Akaike information criterion (AIC) and average posterior probability (APP) were used as criteria to confirm the goodness of fit. A cubic regression was applied. The trajectory analysis was conducted on two age-groups ≤50 and >50 years as previous studies have suggested that changes in BMI and physical activity may vary depending on the age<sup>19 20</sup>.

The sensitivity analysis was conducted by dividing both age groups by gender. No adjustments for co-variables were made.

1  
2  
3 168 All the analyses were performed using Stata/IC Statistical Software: Release 16. College  
4  
5 169 Station (StataCorp LP, TX, USA). The additional Stata module 'traj' was required to conduct group-  
6  
7 170 based trajectory analysis. The module is freely available for both SAS® and Stata software (Jones and  
8  
9 171 Nagin 1999; 2013).

11  
12 172 **Patient and public involvement**

13  
14 173 Participants of research were not involved in setting the study question and outcome measures and  
15  
16 174 were not involved in the design and implementation of the study or writing the manuscript.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**RESULTS**

During the 16-year follow-up, the 66,852 participants had reported body weight and height on average in 3.5 (SD 1.3) study waves and physical activity in 3.6 (SD 1.3) study waves. The sample was predominated by 53,468 women (80%). In the younger group ( $\leq 50$ -year-olds) mean age was 39.8 (SD 7.2), BMI at baseline was 24.6 (SD 4.0)  $\text{kg/m}^2$  and average physical activity was 28.8. (SD 25.5) MET-h/week. In the older group ( $>50$ -year-olds), age was 55.0 (SD 2.9), BMI 25.6 (SD 4.2)  $\text{kg/m}^2$  and physical activity 26.7 (SD 24.1) MET-h/week.

A four-trajectory model was chosen as the five-trajectory model had resulted in a smallest group below a pre-agreed cut-off of 5% (Table 1). Four concurrent trajectories of BMI and physical activity were identified for both age groups (Figure 1 and Figure 2):

- Group 1 (38% among  $\leq 50$  years, 32% among  $> 50$  years): Individuals with normal weight (BMI  $<25 \text{ kg/m}^2$ ) and high level of physical activity (30-35 MET-h/week).
- Group 2 (39% among  $\leq 50$  years, 42% among  $> 50$  years): Individuals with overweight (BMI 25 to 30  $\text{kg/m}^2$ ) and moderately high level of physical activity (25-30 MET-h/week).
- Group 3 (18% among  $\leq 50$  years, 21% among  $> 50$  years): Individuals with obesity (BMI 30 to 35  $\text{kg/m}^2$ ) and moderately low level of physical activity (20-25 MET-h/week).
- Group 4 (5% among  $\leq 50$  years, 5% among  $> 50$  years): Individuals with severe obesity (BMI  $>35 \text{ kg/m}^2$ ) and low level of physical activity ( $<20$  MET-h/week).

**Group 1: Individuals with normal weight and high level of physical activity**

In this group, the younger respondents demonstrated a stable high level of physical activity with a slight rise towards the end of follow-up and their BMI increased slightly throughout the follow-up. For the older respondents, the level of physical activity decreased markedly during the follow-up, even if there was a slight rising pattern in the middle of follow-up. At the same time, the trajectory of BMI remained flat.

**Group 2: Individuals with overweight and moderately high level of physical activity**

201 In this group, the level of physical activity declined in both age groups, but the decline was steeper  
202 among the older respondents. In younger respondents, the decrease of physical activity slowed down  
203 slightly towards the end of follow-up. Simultaneously, BMI was steadily growing among younger  
204 respondents, while remaining relatively flat in older group.

205 **Group 3: Individuals with obesity and moderately low level of physical activity**

206 The physical activity and BMI trajectories were similar to the trajectories observed in group of  
207 overweight individuals with moderately high level of physical activity (group #2), but with a slightly  
208 steeper decline in physical activity and steeper increase in BMI.

209 **Group 4: Individuals with severe obesity and low level of physical activity**

210 Also in this group, physical activity decreased and BMI increased. In younger respondents, this  
211 development slowed down at the end follow-up for both physical activity and BMI. Instead, in older  
212 respondents, the decrease in physical activity accelerated towards the end of follow-up with  
213 simultaneous slight decline in BMI.

214 **Sensitivity analysis**

215 Stratifying the respondents by gender in addition to age resulted in similar findings with few  
216 exceptions (supplementary figures E1-E4 and supplementary table E2). Among normal weight or  
217 overweight respondents, the decline in physical activity was steeper among men compared to women.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**DISCUSSION**

This prospective cohort study in 66,852 public sector employees followed repeatedly by 4-year intervals investigated trajectories of concurrent changes in BMI and physical activity over 16 years. Four trajectory clusters were identified for both participants aged  $\leq 50$  and for those  $> 50$  years: 1) individuals with normal weight and high level of physical activity; 2) individuals with overweight and moderately high level of physical activity; 3) individuals with obesity and moderately low level of physical activity; and 4) individuals with severe obesity and low level of physical activity. On average, BMI increased and physical activity decreased during the follow-up. Some trajectories demonstrated, however, distinctive features. Over time, the respondents with normal weight or overweight gained only a little weight while preserved a high or moderately high level of physical activity, even if the intensity of physical activity mildly decreased especially in older respondents. The decrease in physical activity and increase in BMI were steeper among the respondents with obesity or severe obesity, who had moderately low or low level of physical activity already at the start of the follow-up. Among the normal weight or overweight respondents, decline in physical activity was steeper among men compared to women.

The observed age-related weight gain is in line with previous studies<sup>4-6,21</sup>, as well as the decline in physical activity<sup>4,22,23</sup>. Previous studies have also shown that an increase in BMI slows down with advancing age, and this was also supported by the present findings – the rise in BMI was steeper in the younger respondents<sup>24,25</sup>. During the follow-up, the decline in physical activity mirrored the increase in BMI. Similar findings have been reported before – several studies conducted among middle-aged adults have observed an association between physical activity and weight gain<sup>10,11,26,27</sup>. This association has been described to be dose-dependent – physically active individuals gain less weight than inactive peers.<sup>11</sup> Current results support this finding, since the increase in BMI was less steep in the more active groups. The amount of activity needed to prevent weight gain has been debated. Some studies have concluded that current activity recommendations are not sufficient enough to prevent weight gain and that there is a need for higher activity to remain in the normally

weighted category<sup>10 11 26</sup>. This is in line with the current findings – only high physical activity was associated with normal weight.

The strengths of the study were long follow-up of 16 years, repeated measurements on physical activity and BMI, and a large sample size. For our knowledge, there are no previous multi-trajectory analyses of the relation between physical activity and BMI conducted in adults.

The study has also some limitations. Physical activity was self-reported and only leisure-time and commuting activity were inquired. Thus, physical activity at work was not considered. The distribution of physical activity intensity was skewed – most of the participants were at least somewhat active, and even in the least active group the mean activity level was approximately 18 MET-h/week, which is approximately the equivalent of three hours of brisk walking weekly. BMI was also based on self-reported weight and height, which may cause recall and information bias, possibly resulting in under-reporting of body weight<sup>28</sup>. Most of the participants had BMI above 25 indicating overweight or obesity (62% in the age group of  $\leq 50$  years and 68% in the older), which may reflect the current overweight and obesity pandemic. The cohort included predominantly working-age women employed in public sector. Therefore, the results might not be directly reflected on the entire population, since there might be variation in behavior, for instance among unemployed people or entrepreneurs. Moreover, a public sector often employs people with higher socioeconomic status, who might have more knowledge and financial resources to healthy lifestyle choices compared to manual workers.

The results may be of interest for both clinicians and other stakeholders with respect to informing measures targeting increasing physical activity and controlling weight, especially among middle-aged people. Additionally, the information on the established trajectories may give people more motivation to change their health behavior. Further research may reveal risk factors that affect developmental trajectories seen in this study. Such factors may be, for example, gender, socioeconomic status, smoking, alcohol consumption and concurrent health disorders among others.

## Conclusions

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

270 Changes in BMI and physical activity might be interconnected. The normal weight or overweight  
271 respondents gained only a little weight while preserved a high or moderately high level of physical  
272 activity. Compared to normal weight trajectories, the decrease of physical activity and increase in BMI  
273 were markedly steeper among the obese or severely obese trajectories, who also had moderately low  
274 or low level of physical activity. The findings were similar for both age groups. Among the normal  
275 weight and overweight trajectories, decline in physical activity was steeper among men compared to  
276 women. Since physical inactivity and overweight are both risk factors for many diseases, more  
277 research is needed to develop interventions that could simultaneously affect both.

For peer review only

## 278 LIST OF ABBREVIATIONS

279 BMI: Body Mass Index

280 FPS: Finnish Public Sector cohort study

281 MET: Metabolic Equivalent of Task

## 283 FIGURE LEGENDS

284 Figure 1. Trajectories of physical activity and body mass index amongst respondents  $\leq 50$  years

285 95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in  
286 the figure. Time between responses is approximately four years.

288 Figure 2. Trajectories of physical activity and body mass index amongst respondents  $> 50$  years

289 95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in  
290 the figure. Time between responses is approximately four years.



291 REFERENCES

292 1. Bull F, Goenka S, Lambert V, et al. Physical Activity for the Prevention of Cardiometabolic Disease.  
293 In: rd, Prabhakaran D, Anand S, et al., eds. Cardiovascular, Respiratory, and Related  
294 Disorders. Washington (DC)2017.

295 2. Bigaard J, Frederiksen K, Tjonneland A, et al. Waist circumference and body composition in  
296 relation to all-cause mortality in middle-aged men and women. *Int J Obes (Lond)*  
297 2005;29(7):778-84. doi: 10.1038/sj.ijo.0802976 [published Online First: 2005/05/27]

298 3. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World*  
299 *Health Organ Tech Rep Ser* 2000;894:i-xii, 1-253. [published Online First: 2001/03/10]

300 4. Sui X, Zhang J, Lee DC, et al. Physical activity/fitness peaks during perimenopause and BMI change  
301 patterns are not associated with baseline activity/fitness in women: a longitudinal study with  
302 a median 7-year follow-up. *Br J Sports Med* 2013;47(2):77-82. doi: 10.1136/bjsports-2011-  
303 090888 [published Online First: 2012/07/10]

304 5. Droyvold WB, Holmen J, Midthjell K, et al. BMI change and leisure time physical activity (LTPA): an  
305 11-y follow-up study in apparently healthy men aged 20-69 y with normal weight at  
306 baseline. *Int J Obes Relat Metab Disord* 2004;28(3):410-7. doi: 10.1038/sj.ijo.0802569  
307 [published Online First: 2004/01/16]

308 6. Rosenberger PH, Ning Y, Brandt C, et al. BMI trajectory groups in veterans of the Iraq and  
309 Afghanistan wars. *Prev Med* 2011;53(3):149-54. doi: 10.1016/j.ypmed.2011.07.001  
310 [published Online First: 2011/07/21]

311 7. Fitzpatrick SL, Rosales AG, Brown SD, et al. Behavioural and psychosocial factors associated with 5-  
312 year weight trajectories within the PORTAL Overweight/Obesity Cohort. *Obes Sci Pract*  
313 2020;6(3):272-81. doi: 10.1002/osp4.411 [published Online First: 2020/06/12]

314 8. Barnett TA, Gauvin L, Craig CL, et al. Distinct trajectories of leisure time physical activity and  
315 predictors of trajectory class membership: a 22 year cohort study. *Int J Behav Nutr Phys Act*  
316 2008;5:57. doi: 10.1186/1479-5868-5-57 [published Online First: 2008/11/08]

317 9. Imes CC, Zheng Y, Mendez DD, et al. Group-Based Trajectory Analysis of Physical Activity Change  
318 in a US Weight Loss Intervention. *J Phys Act Health* 2018;15(11):840-46. doi:  
319 10.1123/jpah.2017-0484 [published Online First: 2018/10/14]

320 10. Jakicic JM, Powell KE, Campbell WW, et al. Physical Activity and the Prevention of Weight Gain in  
321 Adults: A Systematic Review. *Med Sci Sports Exerc* 2019;51(6):1262-69. doi:  
322 10.1249/MSS.0000000000001938 [published Online First: 2019/05/17]

323 11. Moholdt T, Wisloff U, Lydersen S, et al. Current physical activity guidelines for health are  
324 insufficient to mitigate long-term weight gain: more data in the fitness versus fatness debate  
325 (The HUNT study, Norway). *Br J Sports Med* 2014;48(20):1489-96. doi: 10.1136/bjsports-  
326 2014-093416 [published Online First: 2014/05/02]

327 12. Kouvonen A, Kivimaki M, Elovainio M, et al. Effort/reward imbalance and sedentary lifestyle: an  
328 observational study in a large occupational cohort. *Occup Environ Med* 2006;63(6):422-7.  
329 doi: 10.1136/oem.2005.020974 [published Online First: 2006/02/25]

330 13. Kouvonen A, Kivimaki M, Virtanen M, et al. Effort-reward imbalance at work and the co-  
331 occurrence of lifestyle risk factors: cross-sectional survey in a sample of 36,127 public sector  
332 employees. *BMC Public Health* 2006;6:24. doi: 10.1186/1471-2458-6-24 [published Online  
333 First: 2006/02/09]

334 14. Leskinen T, Stenholm S, Aalto V, et al. Physical activity level as a predictor of healthy and chronic  
335 disease-free life expectancy between ages 50 and 75. *Age Ageing* 2018;47(3):423-29. doi:  
336 10.1093/ageing/afy016 [published Online First: 2018/03/17]

337 15. Kujala UM, Kaprio J, Sarna S, et al. Relationship of leisure-time physical activity and mortality: the  
338 Finnish twin cohort. *JAMA* 1998;279(6):440-4. doi: 10.1001/jama.279.6.440 [published  
339 Online First: 1998/02/18]

16. Ainsworth BE, Haskell WL, Leon AS, et al. Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc* 1993;25(1):71-80. doi: 10.1249/00005768-199301000-00011 [published Online First: 1993/01/01]
17. Tanasescu M, Leitzmann MF, Rimm EB, et al. Exercise type and intensity in relation to coronary heart disease in men. *JAMA* 2002;288(16):1994-2000. doi: 10.1001/jama.288.16.1994 [published Online First: 2002/10/22]
18. Fogelholm M, Suni, J., Rinne, M., Oja, P., Vuori, I. Physical Activity Pie: A Graphical Presentation Integrating Recommendations for Fitness and Health. *Journal of Physical Activity and Health* 2005;2:391-96.
19. Stenholm S, Pulakka A, Kawachi I, et al. Changes in physical activity during transition to retirement: a cohort study. *Int J Behav Nutr Phys Act* 2016;13:51. doi: 10.1186/s12966-016-0375-9 [published Online First: 2016/04/17]
20. Stenholm S, Solovieva S, Viikari-Juntura E, et al. Change in body mass index during transition to statutory retirement: an occupational cohort study. *Int J Behav Nutr Phys Act* 2017;14(1):85. doi: 10.1186/s12966-017-0539-2 [published Online First: 2017/06/28]
21. Yang Y, Dugue PA, Lynch BM, et al. Trajectories of body mass index in adulthood and all-cause and cause-specific mortality in the Melbourne Collaborative Cohort Study. *BMJ Open* 2019;9(8):e030078. doi: 10.1136/bmjopen-2019-030078 [published Online First: 2019/08/12]
22. Achttien RJ, van Lieshout J, Wensing M, et al. The decline in physical activity in aging people is not modified by gender or the presence of cardiovascular disease. *Eur J Public Health* 2020;30(2):333-39. doi: 10.1093/eurpub/ckz159 [published Online First: 2019/09/29]
23. McAuley E, Hall KS, Motl RW, et al. Trajectory of declines in physical activity in community-dwelling older women: social cognitive influences. *J Gerontol B Psychol Sci Soc Sci* 2009;64(5):543-50. doi: 10.1093/geronb/gbp049 [published Online First: 2009/06/17]
24. Brown WJ, Kabir E, Clark BK, et al. Maintaining a Healthy BMI: Data From a 16-Year Study of Young Australian Women. *Am J Prev Med* 2016;51(6):e165-e78. doi: 10.1016/j.amepre.2016.09.007 [published Online First: 2016/11/22]
25. Lewis CE, Jacobs DR, Jr., McCreath H, et al. Weight gain continues in the 1990s: 10-year trends in weight and overweight from the CARDIA study. Coronary Artery Risk Development in Young Adults. *Am J Epidemiol* 2000;151(12):1172-81. doi: 10.1093/oxfordjournals.aje.a010167 [published Online First: 2000/07/25]
26. Erlichman J, Kerbey AL, James WP. Physical activity and its impact on health outcomes. Paper 2: Prevention of unhealthy weight gain and obesity by physical activity: an analysis of the evidence. *Obes Rev* 2002;3(4):273-87. doi: 10.1046/j.1467-789x.2002.00078.x [published Online First: 2002/12/03]
27. Hankinson AL, Daviglus ML, Bouchard C, et al. Maintaining a high physical activity level over 20 years and weight gain. *JAMA* 2010;304(23):2603-10. doi: 10.1001/jama.2010.1843 [published Online First: 2010/12/16]
28. Connor Gorber S, Tremblay M, Moher D, et al. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev* 2007;8(4):307-26. doi: 10.1111/j.1467-789X.2007.00347.x [published Online First: 2007/06/21]









1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


TABLES


Table 1. Goodness of fit of group-based trajectory analysis models. The chosen models are shown in bold.

Model	Smallest group		BIC <sup>1</sup>	AIC <sup>2</sup>	APP <sup>3</sup>
	n	%			
≤50 years					
1-cluster model	31,797	100%	-905,561	-905,509	1
2-cluster model	8,234	26%	-869,531	-869,432	0.94
3-cluster model	3,331	10%	-851,542	-851,397	0.92
<b>4-cluster model</b>	<b>1,490</b>	<b>5%</b>	<b>-841,703</b>	<b>-841,510</b>	<b>0.89</b>
5-cluster model	898	3%	-835,396	-835,157	0.87
>50 years					
1-cluster model	35,055	100%	-869,200	-869,148	1
2-cluster model	9,690	28%	-836,174	-836,076	0.93
3-cluster model	3,845	11%	-819,600	-819,454	0.91
<b>4-cluster model</b>	<b>1,888</b>	<b>5%</b>	<b>-809,601</b>	<b>-809,409</b>	<b>0.89</b>
5-cluster model	999	3%	-803,977	-803,738	0.87

<sup>1</sup> BIC = Bayesian Information Criterion, <sup>2</sup> AIC = Akaike information criterion, <sup>3</sup> APP = Smallest average posterior probability







For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

Figure E1. Trajectories of physical activity and body mass index (BMI) amongst men < 50 years

95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.




Figure E2. Trajectories of physical activity and body mass index amongst men > 50 years  
95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.






Figure E3. Trajectories of physical activity and body mass index amongst women < 50 years

95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.






Figure E4. Trajectories of physical activity and body mass index amongst women > 50 years  
95% confidence limits are shown as dot-lines. For BMI, very narrow 95% CIs are poorly separable in the figure. Time between responses is approximately four years.

Table E1. Defining the level of physical activity in the survey.

What was the intensity and frequency of your average physical activity (leisure or commuting) during the past year (or since the onset of your disease if the disease had begun less than a year ago)?					
Intensity (Mark all four options)	Amount per week				
	None	<½ hour	1 hour	2 – 3 hours	≥ 4 hours
Normal walking or respective					
Brisk walking or respective					
Light jogging or respective					
Brisk jogging or respective					
The responses were converted into MET units according to a following scheme.					
Intensity (Mark all four options)	MET minutes per week				
	None	<½ hour	1 hour	2 – 3 hours	≥ 4 hours
Normal walking or respective	0	69	138	345	550
Brisk walking or respective	0	99	198	495	792
Light jogging or respective	0	210	420	1050	1680
Brisk jogging or respective	0	240	480	1200	1920

Table E2. Goodness of fit of group-based trajectory analysis models. The chosen models are shown in bold.

Model	Smallest group		BIC <sup>1</sup>	AIC <sup>2</sup>	APP <sup>3</sup>
	n	%			
Men <51 years					
1-cluster	5,894	100%	-156,412	-156,369	1
2-cluster	1,469	25%	-151,020	-150,938	0.93
3-cluster	509	9%	-148,201	-148,080	0.91
<b>4-cluster</b>	<b>292</b>	<b>5%</b>	<b>-146,715</b>	<b>-146,555</b>	<b>0.88</b>
5-cluster	147	2%	-145,799	-145,600	0.86
Men >50 years					
1-cluster	7,490	100%	-177,574	-177,530	1
2-cluster	1,894	25%	-171,451	-171,368	0.92
3-cluster	622	8%	-168,332	-168,209	0.90
<b>4-cluster</b>	<b>334</b>	<b>4%</b>	<b>-166,442</b>	<b>-166,280</b>	<b>0.88</b>
5-cluster	174	2%	-165,267	-165,066	0.87
Women <51 years					
1-cluster	25,903	100%	-746,837	-746,786	1
2-cluster	6,530	25%	-715,572	-715,475	0.95
3-cluster	2,773	11%	-700,393	-700,250	0.92
<b>4-cluster</b>	<b>1,173</b>	<b>5%</b>	<b>-692,029</b>	<b>-691,840</b>	<b>0.90</b>
5-cluster	745	3%	-686,684	-686,449	0.87
Women >50 years					
1-cluster	27,565	100%	-690,012	-689,961	1
2-cluster	7,608	28%	-662,602	-662,506	0.94
3-cluster	3,164	11%	-649,085	-648,944	0.91
<b>4-cluster</b>	<b>1,536</b>	<b>6%</b>	<b>-641,136</b>	<b>-640,949</b>	<b>0.89</b>
5-cluster	842	3%	-636,666	-636,433	0.86

<sup>1</sup> BIC = Bayesian Information Criterion, <sup>2</sup> AIC = Akaike information criterion, <sup>3</sup> APP = Smallest average posterior probability

STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page No
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	4
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	6,7
Objectives	3	State specific objectives, including any prespecified hypotheses	7
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	8
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (b) For matched studies, give matching criteria and number of exposed and unexposed	8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	8
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	8
Bias	9	Describe any efforts to address potential sources of bias	9
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	9
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, explain how loss to follow-up was addressed (e) Describe any sensitivity analyses	9 9 9 9 9
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	10
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) Summarise follow-up time (eg, average and total amount)	10
Outcome data	15*	Report numbers of outcome events or summary measures over time	10,11

1	Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	10,11
2			(b) Report category boundaries when continuous variables were categorized	
3			(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
4				
5				
6				
7				
8				
9	Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10,11
10				
11	<b>Discussion</b>			
12				
13	Key results	18	Summarise key results with reference to study objectives	12
14	Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	13
15				
16				
17	Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	12,13
18				
19	Generalisability	21	Discuss the generalisability (external validity) of the study results	13
20				
21	<b>Other information</b>			
22	Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	2
23				
24				

25  
26 \*Give information separately for exposed and unexposed groups.

27  
28 **Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and  
29 published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely  
30 available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at  
31 <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is  
32 available at <http://www.strobe-statement.org>.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60